



## Adult Grand Rounds

Mike R. Schoenberg, PhD, ABPP – Chair

Gregory Lee, PhD, ABPP – Discussant

Clea Evans, PhD - Discussant

Robin Hilsabeck, PhD, ABPP - Discussant



## FINANCIAL DISCLOSURE

### M. Schoenberg

- Financial support from State of FL, USF Practice plan, medicolegal consulting (< 30% of annual salary), Oxford University Press, Springer, Research support from AACN Foundation and U. of South Florida MCOM

### C. Evans

- None to report

### G. Lee

- Financial support from the State of Georgia, MCG Practice Plan, National Academy of Neuropsychology, Oxford University Press, Taylor & Francis, Veloxis Pharmaceuticals A/S

### R. Hilsabeck

- Royalties from Oxford University Press and John Wiley & Sons.



## The man who used to know everything

Grand Rounds - Adult  
National Academy of Neuropsychology 38<sup>th</sup> Annual Conference  
October 17<sup>th</sup>, 2018

Justin B. Miller, Ph.D., ABPP/CN  
Cleveland Clinic Lou Ruvo Center for Brain Health  
Las Vegas, Nevada



## Financial Disclosures

- I have **no relevant** financial relationships to disclose:
- Employee of: Cleveland Clinic
- Consultant for: Philips HealthWorks
- Stockholder in: None
- Research support from: Avanir Pharmaceuticals
- Honoraria from: None



## Demographics

- 47 year old, right handed, Filipino man
- Born in Germany, raised throughout the world (U.S. Military Family)
- 12 years of education; completed in English
  - Academic history largely known
- Married once, two children (ages 4 and 6)
- Employed full-time in shipping/receiving; worked the docks
  - Released from employment 4 weeks prior to present evaluation secondary to declining performance
  - On FMLA at time of eval



## Referral Question

- Presented to outpatient neurology center, specializing in neurodegenerative disease
  - Primary complaints: short and long term memory loss
- Evaluated by cognitive disorders neurologist
  - Severe impairment noted on cognitive screening (MoCA = 7/30)
  - Neurological exam was unremarkable
- Patient was clinically referred for differential diagnosis and treatment planning
  - Question about need for permanent disability had been raised



## Relevant History

- September 2016:
  - Patient has 2<sup>nd</sup> heart attack (first in 2004); multiple stents placed without complications
  - Was reportedly doing okay after hospitalization
- December 2016:
  - Patient's wife begins noticing that he is having difficulty completing routine tasks around the home (e.g., forgetting items while grocery shopping)
- July 2017:
  - Patient's memory declined markedly
  - Difficulty recalling names of family members (inc. children, wife)
  - Still working, but employer had to make changes to accommodate patient
  - Started becoming less engaged with family members
  - Had been fully independent in all basic and complex ADLs



## Relevant History – Hospitalization

- July 2017
  - Emergently admitted to hospital by PCP
    - Hospital workup: noncontrast MRI, MRA, EEG
    - Only finding was a small posterior fossa cyst (18mm); all studies otherwise normal
  - Attending physician questioned whether Brilinta was the cause
    - Only treatment change was to replace with Brilinta with clopidogrel
    - Discharged home in stable condition, without supports



## Relevant History

- July 2017 – March 2018:
  - Patient continued to decline cognitively
  - Lost his job as a laborer; on FLMA and determining the need for permanent disability
- At time of eval (per wife):
  - Starting to get "pieces" of his memory back
  - Not asking repetitive questions as often
  - Has been more outgoing, like his old self
  - Resumed speaking with cousins and other family members
  - Able to go to the store with a list and get all items



## Presenting Complaints – March 2018

- Majority of his long term and autobiographical memory are gone
  - Unable to state his mother's name, where he was born and raised, the name of his last employer, the name of his children's school, where his wife works, or any recent president.
- Remembers daily schedule without error
  - Dropping his wife off at work
  - Taking his children to and from school
  - Does not get lost while driving
- Recognizes faces readily (though may not always remember the name)
- Significant problems with auditory comprehension
  - Often requires simplification and repetition to ensure he understands
  - No reports of expressive language difficulty



## Presenting Complaints – March 2018

- Patient stated that he is "sad at myself" when asked about his mood
- Acknowledged getting easily frustrated and his wife noted that he has seemed a bit more moody lately, though this is improving.
- Problems with significant apathy; requires prompting to engage in most tasks, though there is no clear anhedonia.
- Suicidal ideation, delusions, and hallucinations were denied.
- Eating less and reports that food does not taste good
  - Lost a considerable amount of weight in the past year
  - No apparent cravings or changes in food preferences.
- Sleep problems denied
  - No reports of REM sleep behavior disorder



## Additional History

- Past Medical History
  - 1<sup>st</sup> Heart attack (2004)
  - 2<sup>nd</sup> Heart attack with multiple coronary stents placed (2016)
    - No known complications
  - Hypertension
  - High cholesterol
  - Coronary artery disease
  - Obstructive sleep apnea (treated)
  - Diabetes mellitus (most recent A1c = 5.7)
- Past Psychiatric history: none
- Family History: late onset of unspecified dementia (paternal grandmother and 2 paternal uncles)



## Current Medications

- Aspirin (low dose)
- Levocetirizine (5mg/day)
- Atorvastatin (20mg/day)
- Clopidogrel (75mg/day)
- Metformin (2000mg/day)
- Losartan (50mg/day)
- Fluticasone



## Preliminary Hypotheses

- Anoxic injury (heart attack; surgical complication)
- Stroke
- Paraneoplastic syndrome
- Brain tumor
- Malingering
- Medication side effects
- Toxic exposure
- Prion disease



## Test battery

- Boston Naming Test
- Delis-Kaplan Executive Functioning System
  - Color Word Interference
  - Trail Making
  - Verbal Fluency
- Hopkins Verbal Learning Test, Revised
- Brief Visuospatial Memory Test, Revised
- Wechsler Memory Scale, 4<sup>th</sup> Ed. – Logical Memory
- Judgment of Line Orientation
- Wechsler Adult Intelligence Scale, 4<sup>th</sup> Ed.
  - Block Design
  - Visual Puzzles
  - Coding
  - Digit Span
  - Similarities
- Test of Practical Judgment
- Wide Range Achievement Test, 4<sup>th</sup> Ed. – Reading
- Medical Symptom Validity Test
- Test of Memory Malingering
- Geriatric Depression Scale



## Behavioral Observations – Interview

- Alert and oriented to person, day, and month
  - Able to identify the date and year with a logical cue
  - Unable to identify the city, state, or name of the clinic even with multiple choice cues
- Affect was very flat and restricted, but pleasant with examiner
- No apparent speech or language errors present
- Speech was soft and slow, but normal rhythm, tone, and prosody
- Expressive language was simple with very short utterances and use of repetitive stock phrases
  - (e.g., referring to his wife as “the love of my life” and “I used to know everything”)
- Thinking was very concrete and literal
  - Only responded to direct questions and had significant difficulty answering
  - No indication of delusion or disorganization.



## Behavioral Observations – Testing

- Easily confused with test instructions and required significant elaboration, simplification, demonstration, and repetition
- Worked slowly but consistently; no evidence of distractibility or disinhibition
- Appeared to stare blankly at test materials without responding but when prompted, he defensively noted that he was trying to work through the item
- Unable to comprehend hypothetical situations intended to elicit judgment and decision making (TOPJ), even with elaboration
- Unable to complete emotional symptom inventories due to comprehension problems and had difficulty recognizing letters on the WRAT-4 Reading subtest
- No difficulty with TMT
- Unable to name a single animal during verbal fluency (!)
- 2 of 60 items correctly on BNT (!!)



## Results – Validity

- Behavioral Observations
  - Loss of autobiographical memory
  - Contrast between daily functioning and presenting complaints
  - BNT and animal naming scores
  - Trouble with letter sequencing but normal number-letter sequencing
- TOMM:
  - Trial 1 = 46
  - Trial 2 = 49
- MSVT:
  - Immediate Recall = 95
  - Delayed Recall = 95
  - Consistency = 90
  - Paired Associates = 0
  - Free Recall = 15
- Reliable Digit Span = 6



## VALID OR INVALID?

### Results

|  | Raw Score | Standard Score | Prev. Testing | Site | Rating          |
|--|-----------|----------------|---------------|------|-----------------|
| WRAT-4 Reading                                 | 18        | 55             | ---           | ---  | Impaired        |
| <i>Wechsler Memory Scale, 4th Ed.</i>          |           |                |               |      |                 |
| Logical Memory Immediate Recall                | 1         | 1              | ---           | ---  | Impaired        |
| Logical Memory Delayed Recall                  | 1         | 1              | ---           | ---  | Impaired        |
| Logical Memory Recognition                     | 18        | ---            | ---           | ---  | Mildly Impaired |
| <i>Hopkins Verbal Learning Test</i>            |           |                |               |      |                 |
| (T-Score)                                      |           |                |               |      |                 |
| Trial 1 / 2 / 3                                | (1/1/3)   | (19/19/19)     | ---           | ---  | ---             |
| Total Recall                                   | 3         | 19             | ---           | ---  | Impaired        |
| Delayed Recall                                 | 0         | 19             | ---           | ---  | Impaired        |
| Percent Retention                              | 0         | 19             | ---           | ---  | Impaired        |
| Rec. Discrimination (H / FA) ✓                 | (5/9)     | -4             | ---           | ---  | Impaired        |
| <i>Brief Visuospatial Memory Test, Revised</i> |           |                |               |      |                 |
| (T-Score)                                      |           |                |               |      |                 |
| Trial 1 / 2 / 3                                | (1/1/3)   | (25/19/19)     | ---           | ---  | ---             |
| Total Recall                                   | 5         | 19             | ---           | ---  | Impaired        |
| Delayed Recall                                 | 1         | 19             | ---           | ---  | Impaired        |
| Percent Retained                               | 33        | ---            | ---           | ---  | Impaired        |
| Discrimination Index (H / FA) ✓                | (5/0)     | 5              | ---           | ---  | Low Average     |



## Results

| Attention / Working Memory / Processing Speed |         |          |    |     |             |             |
|---|---------|----------|----|-----|-------------|-------------|
| <i>D-KEFS Trail Making Test</i>               |         |          |    |     |             |             |
| Scanning                                      | 0       | 35       | 5  | --- | 5 Mild Imp. |             |
| Numbers                                       | 0       | 50       | 6  | --- | 9 Low Avg.  |             |
| Letters                                       | 0       | 59       | 3  | --- | 1 Impaired  |             |
| Speed   | 0       | 34       | 10 | --- | 50 Average  |             |
| <i>D-KEFS Color-Word Interference</i>         |         |          |    |     |             |             |
| Color Naming                                  | Errors: | Sut, 2sc | 90 | 1   | ---         | <1 Impaired |
| Word Reading                                  | Errors: | Dut, 1sc | 68 | 1   | ---         | <1 Impaired |
| WAIS-IV Coding                                |         |          | 44 | 6   | ---         | 9 Low Avg.  |
| WAIS-IV Digit Span Total                      |         |          | 15 | 4   | ---         | 2 Mild Imp. |
| Forward                                       | LDSF:   |          | 4  | 6   | ---         | 5 Mild Imp. |
| Backward                                      | LDSB:   |          | 3  | 5   | ---         | 9 Low Avg.  |
| Sequencing                                    | LDS:    |          | 3  | 4   | ---         | 5 Mild Imp. |

## Results

| Languages  |         |          |     |     |                  |             |
|--|---------|----------|-----|-----|------------------|-------------|
| Boston Naming Test                               | 2       | T = 9    | --- | --- | ---              |             |
| Visual Perception/Organization                   |         |          |     |     |                  |             |
| BVMT - Copy                                      | 10      | ---      | --- | --- | WNI              |             |
| Judgment of Line Orientation                     | 23      | ---      | --- | --- | Impaired         |             |
| WAIS-IV Block Design                             | 24      | 6        | --- | --- | 9 Low Avg.       |             |
| WAIS-IV Visual Puzzles                           | 7       | 6        | --- | --- | 9 Low Avg.       |             |
| Executive Functioning                            |         |          |     |     |                  |             |
| WAIS-IV Similarities                             | 0       | 1        | --- | --- | <1 Impaired      |             |
| <i>Delis-Kaplan Executive Functioning System</i> |         |          |     |     |                  |             |
| (4/7/4)  |         |          |     |     |                  |             |
| Phonemic Fluency (FAS)                           | 15      | 3        | --- | --- | -2.65 1 Impaired |             |
| Animals  | 0       | ---      | --- | --- | -4.06 ---        |             |
| Category Fluency (Animals, Boys names)           | 5       | 1        | --- | --- | <1 Impaired      |             |
| Category Switching                               | 0       | 1        | --- | --- | <1 Impaired      |             |
| CW Inhibition                                    | Errors: | 2uc, 0sc | 180 | 1   | ---              | <1 Impaired |
| CW Switching                                     | Errors: | 1uc, 0sc | 177 | 1   | ---              | <1 Impaired |
| Trails Switching                                 | Errors: | 0        | 98  | 8   | ---              | 25 Average  |



## VALID OR INVALID?

## CLINICAL IMPRESSIONS



### Confirmatory bias

- The tendency to search for, interpret, favor, and recall information in a way that confirms one's preexisting beliefs or hypotheses (Plous, 1999)
  - Selective gathering of supporting info and neglect of contradictory info
  - Particularly prevalent in forensic settings (Deja-Murti and Loshart, 2012)
- Avoid by intentionally seeking out disconfirming evidence
  - E.g., are there other reasons why this person might have done so badly?
- Multidisciplinary clinics may be less vulnerable to these biases because of the wealth of additional information available
  - Updated neuroimaging drastically changed clinical impressions




### Evidence of malingering

- |  |   |
|--|---|
| <p><b>Supporting</b></p> <ul style="list-style-type: none"> <li>Very young for almost all neurodegenerative disease</li> <li>Loss of autobiographical information unusual, even for severe dementia</li> <li>Level of cognitive impairment highly discordant with functioning</li> <li>Normal neurological exam</li> <li>Historical imaging reported normal</li> <li>Disability claim pending</li> </ul> | <p><b>Disconfirming</b></p> <ul style="list-style-type: none"> <li>Passed all validity indices                             <ul style="list-style-type: none"> <li>Could he have been coached?</li> </ul> </li> <li>Expressive language errors (i.e., repetitive stock phrases)</li> <li>Evidence of poor comprehension</li> </ul> |
|--|---|




### Disconfirming Evidence – Brain MRI

- No evidence of an acute intracranial process or intracranial mass.
- Moderate generalized volume loss by visual inspection.
- Hippocampal volumes at the 1st percentile when compared to age matched normal controls by quantitative analysis
  - Severe hippocampal formation volume loss by visual inspection.
- Minimal white matter disease which is nonspecific but likely reflective of chronic microvascular ischemia.



### FDG-PET

- Diffuse cortical hypometabolism, more pronounced in association cortex in bilateral temporal, left parietal, occipital and frontal regions
- Distribution pattern suggests neurodegenerative disorders with differentials including diffuse Lewy-body dementia
- Other likely differentials include temporal dominant atypical Alzheimer's dementia, frontotemporal dementia



### CLINICAL IMPRESSIONS



### Clinical Diagnosis – Semantic dementia

|   |  |
|---|--|
| <p><b>1. Clinical diagnosis of semantic variant PPA</b></p> <p>Both of the following core features must be present:</p> <ol style="list-style-type: none"> <li>Impaired confrontation naming</li> <li>Impaired single word comprehension</li> </ol> <p>At least 2 of the following other diagnostic features must be present:</p> <ol style="list-style-type: none"> <li>Impaired object knowledge, particularly for low frequency or low familiarity items</li> <li>Surface dyslexia or dysgraphia</li> <li>Spelling impairment</li> <li>Spoken speech production (grammar and motor speech)</li> </ol> <p><b>3. Imaging supported semantic variant PPA diagnosis</b></p> <p>Both of the following criteria must be present:</p> <ol style="list-style-type: none"> <li>Clinical diagnosis of semantic variant PPA</li> <li>Imaging must show one or more of the following results:                             <ol style="list-style-type: none"> <li>Frontotemporal atrophy (temporo-parietal)</li> <li>Prefrontal anterior temporal lobe atrophy or hypometabolism on SPECT or PET</li> </ol> </li> </ol> | <ul style="list-style-type: none"> <li>Variant of frontotemporal dementia (FTD)</li> <li>Further subtyped as a primary progressive aphasia (PPA)                             <ul style="list-style-type: none"> <li>Semantic dementia is primarily characterized by loss of semantic knowledge</li> <li>Presents as anomia and single-word comprehension deficits</li> <li>With progression, stock phrases become common</li> </ul> </li> <li>Brain atrophy typically greater on the left</li> <li>Caused by accumulation of TDP-43</li> <li>Typical age of onset: 50s – 60s</li> <li>Limited inheritance</li> </ul> |
|---|--|





### Conclusions

- Confirmatory bias is a common source of error in clinical decision making
  - Must remain vigilant for it and actively seek disconfirming evidence
- Veracity of PVTs:
  - Patient with severe, nearly global impairment
    - Still able to perform above criterion on TOMM and MSVT
  - Patient with severe loss of semantic knowledge
    - Still able to perform above criterion on TOMM and MSVT!
  - Hippocampus almost completely absent (quantified at 1<sup>st</sup> percentile)
    - Still able to perform above criterion on TOMM and MSVT!!



### History of Nonverbal Learning Disability (NLD)

- Study of LD began around 1890 (Spreen, 2011)
- Initially focused on ability to read and write (Orton, 1937)
- “Nonverbal Disorders of Learning” & “Nonverbal Learning Disability” (Johnson & Myklebust, 1967; Myklebust, 1975)
  - Selective deficits in: perception, processing of gestures, motor learning, body image, spatial orientation, and left-right orientation
- Furthered by the work of Byron Rourke (1989 & 1995)
  - Visual-spatial learning
  - Tactile and motor skills
  - Mathematics



### Controversy

- Definition of a learning disability
- Variable diagnostic criteria creating tautological problems
- Methodological issues in the literature

(Fine et al., 2013 & Spreen, 2011)



### Proposed Diagnostic Criteria for NLD

- **Criterion A.** A persistent deficit in one or more measures of nonverbal intelligence or reasoning in the presence of an average or above-average verbal intelligence
- **Criterion B.** Substantial weaknesses, currently or emerging from the child’s history, in processing visuospatial information, as manifested by at least two of the following weaknesses
  1. Difficulties in perceiving organized forms
  2. Difficulties in reproducing simple drawings by copy or memory
  3. Difficulties in temporarily remembering and manipulating visuospatial information


Comolli, C., Mammarella, I. C., & Fine, J. G., 2016



### Proposed Diagnostic Criteria for NLD, Cont.

- **Criterion C.** Presence of clinical and/or psychometric indexes of weaknesses in at least one of the following areas, currently or by history
  1. Fine-motor impairments including praxis and/or output
  2. Poor academic achievement in activities involving visuospatial skills, mathematics, or other, in the presence of an average or above-average performance in reading decoding tasks
  3. Difficulties in social interactions

Comolli, C., Mammarella, I. C., & Fine, J. G. (2016)



38TH ANNUAL CONFERENCE  
OCTOBER 17-20, 2018

BECOMING AGENTS OF CHANGE

SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

### Proposed Diagnostic Criteria for NLD, Cont.

- **Criterion D.** Several symptoms were present before age 7, although they could not have become fully manifest until academic demands exceeded the children's capacities, or were masked by good verbal strategies
- **Criterion E.** There is clear evidence that the symptoms interfere with, or reduce the quality of, academic, occupational, or social functioning
- **Criterion F.** These disorders are not better explained by the presence:
  - ASD
  - DCD
  - ID

Comoldi, C., Mammarella, I. C., & Fine, J. G. (2016)

Samaritan Health Services




38TH ANNUAL CONFERENCE  
OCTOBER 17-20, 2018

BECOMING AGENTS OF CHANGE

SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

### Case Presentation

Samaritan Health Services



38TH ANNUAL CONFERENCE  
OCTOBER 17-20, 2018


BECOMING AGENTS OF CHANGE

SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

### Case Presentation

- **Demographic Data**
  - 25 year-old, single, non-Hispanic White female completing her 16<sup>th</sup> year of education; majoring in English
  - English is her only language
- **Referral**
  - Differential diagnosis of ADHD and LD
  - Accommodations

Samaritan Health Services



38TH ANNUAL CONFERENCE  
OCTOBER 17-20, 2018

BECOMING AGENTS OF CHANGE

SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

### Pertinent Psychosocial/ Educational History

- Normal Pregnancy, met all developmental milestones on time.
- "Average" grades throughout elementary school (public school)
  - Reported multiple reading evaluations
  - Speech therapy in 2<sup>nd</sup>, 3<sup>rd</sup>, & 4<sup>th</sup> grades
- Home schooled for middle school and high school
- Did not take test for high school diploma "because of test anxiety"
- Passed GED exam on the fourth attempt
- Received accommodation in college
  - Extended time for writing
- Reported that she needed additional accommodations for math and reading

Samaritan Health Services



38TH ANNUAL CONFERENCE  
OCTOBER 17-20, 2018


BECOMING AGENTS OF CHANGE

SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

### Pertinent Medical History

- **Headaches**
  - Migraines: 3 per year
  - Tension: 4 per month
- "Hypermobility Syndrome" in hands
- **Sleep**
  - Sleep apnea (on CPAP)
- Unremarkable neurological history

Samaritan Health Services



38TH ANNUAL CONFERENCE  
OCTOBER 17-20, 2018


BECOMING AGENTS OF CHANGE


SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

### Psychosocial History

- **Mood:** "More angry than I should be and extremely stressed out."
- **Lifelong anxiety**
  - Hair pulling
  - Hitting
- Depression and passive SI beginning at 15 years old
- Currently participating in psychotherapy – "Helpful"


Samaritan Health Services




**BECOMING AGENTS OF CHANGE**  
 38TH ANNUAL CONFERENCE  
 OCTOBER 17-20, 2018  
 SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

## Collateral Report (Mother)

- Spoke in full sentences at 18 months and was able to read before starting the first grade
- ADHD Symptoms (since toddlerhood)
  - Missing details/producing inaccurate work
  - Easily distracted
  - Not sustaining attention
  - Starting but not completing work
  - Maintaining a messy environment
  - Losing personal belongings
- Deficits negatively impacted her academic performance, social interaction, and relationships with teachers
- "Good grades" precluded her from being tested for ADHD and LD
- Lifelong Anxiety






**BECOMING AGENTS OF CHANGE**  
 38TH ANNUAL CONFERENCE  
 OCTOBER 17-20, 2018  
 SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

## Chief Complaints

- Attention Deficits**
  - Missing details/producing inaccurate work
  - Difficulty sustaining attention
  - Not listening when spoken to
  - Starting but not finishing work
  - Maintaining a messy/disorganized environment
  - Avoiding tasks that require sustained mental effort
  - Losing personal belongings frequently
  - Becoming easily distracted
- Hyperactivity**
  - Fidgeting/squirming
  - Leaving her seat and running at in appropriate times when younger
  - Difficulty engaging in leisurely activities
  - Feeling down by a mirror
  - Talking excessively
  - Interrupting others
  - Difficulty waiting her turn
- Mathematics**
  - Transposing numbers
  - Making up math signs
  - No improvement following tutoring






**BECOMING AGENTS OF CHANGE**  
 38TH ANNUAL CONFERENCE  
 OCTOBER 17-20, 2018  
 SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

## Test Results








**BECOMING AGENTS OF CHANGE**  
 38TH ANNUAL CONFERENCE  
 OCTOBER 17-20, 2018  
 SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

## Behavioral Observations

- Braces on her hands bilaterally
- Appropriate eye contact
- Adequate interpersonal skills
- Difficulty sitting still
- Easily distracted
- Tangential








**BECOMING AGENTS OF CHANGE**  
 38TH ANNUAL CONFERENCE  
 OCTOBER 17-20, 2018  
 SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA


## Performance Validity

| WMT | %    |
|-----|------|
| IR  | 92.5 |
| DR  | 97.5 |
| CNS | 90   |

|                 | Raw | RDS      | Raw | RDS | Overall Clinical Sample |
|-----------------|-----|----------|-----|-----|-------------------------|
| ACS Word Choice | 50  | Forward  | 5   | 10  | <75%                    |
|                 |     | Backward | 5   |     |                         |







**BECOMING AGENTS OF CHANGE**  
 38TH ANNUAL CONFERENCE  
 OCTOBER 17-20, 2018  
 SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

## Intellectual Functioning

| WAIS-IV          | SS  | Comparison to the Overall Normative Sample |                 |
|------------------|-----|--|-----------------|
|                  |     | Raw  | Cumulative %age |
| VCI              | 150 |  |                 |
| PII              | 77  |  |                 |
| WMI              | 92  | VCI-PII                                    | 73 <-0.2        |
| PSI              | 76  | VCI-WMI                                    | 58 <-0.1        |
| FSIQ             | 102 | VCI-PSI                                    | 74 1            |
|                  | Raw | SS   |                 |
|                  |     | PRI-WMI                                    | -15 13.1        |
| Similarities     | 36  | PRI-PSI                                    | 1 47            |
| Vocabulary       | 55  | WMI-PSI                                    | 16 14.9         |
| Block Design     | 24  |  |                 |
| Matrix Reasoning | 12  |  |                 |
| Digit Span       | 29  |  |                 |
| Arithmetic       | 11  |  |                 |
| Symbol Search    | 23  |  |                 |
| Coding           | 44  |  |                 |









38TH ANNUAL CONFERENCE  
OCTOBER 17-20, 2018

SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA


BECOMING AGENTS OF CHANGE

### Differential Diagnosis: ASD

- Absence of restrictive/repetitive patterns of behavior, interests, or activities
- More severe in ASD
  - Sameness/inflexible adherence to routines
  - Reactivity to sensory input
  - Pragmatic difficulties
    - Unusual prosody
    - Verbose speech
    - Difficulties interpreting jokes / figurative language
    - Lack of response to nonverbal social cues
- Difficulties with visuospatial reasoning not consistent in children with ASD

Samaritan Health Services

Semrud-Clickeman et al., 2010



38TH ANNUAL CONFERENCE  
OCTOBER 17-20, 2018

SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA


BECOMING AGENTS OF CHANGE

### Differential Diagnosis: SLD

- Considerable overlap with mathematics LD
- Geary's (2004) 3-subtypes of mathematics LD
  - Procedural } More likely to be represented by pure mathematics LD
  - Semantic } More likely to be represented in NLD
  - Visuospatial } More likely to be represented in NLD

Samaritan Health Services

Mazzocco & Myers, 2003



38TH ANNUAL CONFERENCE  
OCTOBER 17-20, 2018

SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

BECOMING AGENTS OF CHANGE

### Differential Diagnosis: ADHD

- Similarities:
  - Difficulties in organizing tasks/activities
  - Visual sustained attention tasks
- Differences
  - NLD does well on verbal sustained attention tasks
  - Visuospatial < Verbal intelligence not documented within ADHD

Samaritan Health Services

Semrud-Clickeman et al., 2010



38TH ANNUAL CONFERENCE  
OCTOBER 17-20, 2018

SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

BECOMING AGENTS OF CHANGE

- ✓ **Criterion A.** A persistent deficit in one or more measures of nonverbal intelligence or reasoning in the presence of an average or above-average verbal intelligence
- ✓ **Criterion B.** Substantial weaknesses, currently or emerging from the child's history, in processing visuospatial information, as manifested by at least two of the following weaknesses
  1. Difficulties in perceiving organized forms ✓
  2. Difficulties in reproducing simple drawings by copy or memory
  3. Difficulties in temporarily remembering and manipulating visuospatial information
- ✓ **Criterion C.** Presence of clinical and/or psychometric indexes of weaknesses in at least one of the following areas, currently or by history
  1. Fine-motor impairments including praxis and/or output ✓
  2. Poor academic achievement in activities involving visuospatial skills, mathematics, or other, in the presence of an average or above-average performance in reading decoding tasks ✓
- ✓ **Criterion D.** Several symptoms were present before age 7, although they could not have become fully manifest until academic demands exceeded the children's capacities, or were masked by good verbal strategies
- ✓ **Criterion E.** There is clear evidence that the symptoms interfere with, or reduce the quality of, academic, occupational, or social functioning
- ✓ **Criterion F.** These disorders are not better explained by the presence:
  - ASD
  - DCD
  - ID

Samaritan Health Services



38TH ANNUAL CONFERENCE  
OCTOBER 17-20, 2018

SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

BECOMING AGENTS OF CHANGE

### Treatment/Recommendations

- Psychiatric/social supports
  - Suicide safety plan
  - Medication management for mood
  - Behavioral activation
  - Improved sleep quality through use of sleep hygiene

Samaritan Health Services



38TH ANNUAL CONFERENCE  
OCTOBER 17-20, 2018

SHERATON NEW ORLEANS HOTEL | NEW ORLEANS, LA

BECOMING AGENTS OF CHANGE

### Treatment/Recommendations

- OT evaluation: fine motor dexterity and adaptation to visuospatial/perceptual challenges in day to day life
- Supports in day to day life for NLD, as it affects inattention:
  - Maintain an organizer with a schedule of daily events and reminders for assignments
  - Limit the number of distractions present when working
  - Budget extra time to complete tests and work knowing that you may become distracted
- Academic accommodations
  - Limit courses heavy in mathematic content
  - Participate in class discussions as much as possible to enhance concentration
  - Study with others to reinforce your understanding of the material
  - Always "over-learn" the material
  - Apply the concepts to a "real-life" scenario
  - Pace completion of mathematics homework

Do not move onto more complex math tasks until you have demonstrated mastery over foundational math concepts

Samaritan Health Services



## Questions...



## References

- Cornoldi, C., Mammarella, I. C., & Fine, J. G. (2016). *Nonverbal learning disabilities*. Guilford Publications.
- Fine, J. G., Semrud-Clikeman, M., Bledsoe, J. C., & Musicki, K. A. (2013). A critical review of the literature on NLD as a developmental disorder. *Child Neuropsychology, 19*(2), 190-223.
- Geary, D. C. (2004). Mathematics and learning disabilities. *Journal of learning disabilities, 37*(1), 4-15.
- Johnson, D. J., & Myklebust, H. R. (1967). *Learning Disabilities: Educational Principles and Practices*.
- Mazocco, M. M., & Myers, G. F. (2003). Complexities in identifying and defining mathematics learning disability in the primary school-age years. *Annals of dyslexia, 53*(1), 218-253.
- Myklebust, H. R. (1975). *Nonverbal learning disabilities: Assessment and intervention*.
- Orton, S. T. (1937). *Reading, writing and speech problems in children*.
- Rourke, B. P. (1989). *Nonverbal learning disabilities: The syndrome and the model*. Guilford Press.
- Rourke, B. P. (Ed.). (1995). *Syndrome of nonverbal learning disabilities: Neurodevelopmental manifestations*. Guilford Press.
- Semrud-Clikeman, M., Walkowiak, J., Wilkinson, A., & Christopher, G. (2010). Neuropsychological differences among children with Asperger syndrome, nonverbal learning disabilities, attention deficit disorder, and controls. *Developmental neuropsychology, 35*(5), 582-600.
- Spreen, O. (2011). Nonverbal learning disabilities: A critical review. *Child Neuropsychology, 17*(5), 418-443.



A Case of Progressive Dementia associated with Autoimmune Encephalitis  
 Pamela F. Willis, Ph.D., ABPP-CN  
 Brooke Army Medical Center  
 17 October 2018



## Purpose and Outline

**Purpose:** To present a case of progressive dementia associated with autoimmune encephalitis with N-type voltage gated calcium channel autoantibodies (VGCC abs)

The views expressed herein are those of the author and do not reflect the official policy or position of Brooke Army Medical Center, the U.S. Army Medical Department, the U.S. Army Office of the Surgeon General, the Department of the Army, the Department of the Air Force and Department of Defense or the U.S. Government.



## Autoimmune Encephalitis

- Onset acute or subacute; typically over days or weeks
- Symptoms may include mood or behavioral changes, short-term memory problems, cognitive dysfunction, psychotic symptoms and possible seizures, sometimes catatonic state
- In about 50% of cases MRI normal
- EEG often abnormal for non-specific slow & disorganized brain wave activity
- Best known types are those with antibodies to NMDAR (Anti-N-Methyl-D-Aspartate Receptor) and VGKC (Voltage-gated potassium channel);
- Limbic encephalitis typically affects mesial temporal lobes (hippocampus) and orbitofrontal cortex but also the cerebellum

Tuzun & Dalmau 2007  
 Loughan et al. 2016



## Autoimmune Encephalitis

- Treatment typically consists of steroids, IVIG, plasma exchange
- With early detection, more than 75% of patients with this type of encephalitis recover completely or have only mild neurological sequelae
- Neuropsychological deficits may be more lasting and can include persistent deficits in attention, working memory, episodic memory, and executive functions
- Patients who received immunotherapy during the initial onset had a better cognitive outcome than those who received immunotherapy later

Finke et al. 2012  
 Loughan et al. 2016



## VGCC

- The VGCC's (voltage-gated calcium channels) are found in several cells, to include smooth and skeletal muscle fibers, endocrine cells, and neurons.
- VGCC's are located on the presynaptic membrane of the axon terminal, open by action potential and lead to the entry of calcium ions into the axon terminals
- Calcium influx results in movement of acetylcholine vesicles towards the presynaptic membrane and acetylcholine is released into the synaptic cleft.
- VGCC abs are divided into 5 types: L, P/Q, N, R, and T
- VGCC P/Q type and N type often associated with Lambert-Eaton Myasthenia Syndrome (LEMS) and lung cancer
- Presence of VGCC abs has been associated with both paraneoplastic and nonparaneoplastic cerebellar degeneration
- Bekircan-Kurt et al. 2015

Bekircan-Kurt et al. 2015



- Mayo 2016 study examined a group of 236 patients with VGCC-P/Q type or VGCC-N type
- Autoimmune neurological diagnoses included encephalopathy, ataxia, myelopathy, neuropathy, neuromuscular junction disorder, and myopathy; LEMS mostly diagnosed separately
- Frequencies of diagnoses higher for those with high titer vs. low or medium expressed as nanomoles per liter of serum
- High=1.00 nmol/L or greater N=7
- Medium=0.10-0.99 nmol/L N=79
- Low=0.04-0.10 nmol/L N=150
- Overall positive predictive value was low at only 22%
- Pts with low values had autoimmune neurological diagnoses in 19%, about 20% normal, but 60% another cause, often neurodegenerative or records had insufficient data

Zalewski et al. 2016



- Median age of neurological symptom onset 57 years
- 52% of seropositive patients were women
- Cognitive disorders occurred in 41/236 (17%) and included degenerative dementia, autoimmune encephalopathy, mild cognitive impairment, and cognitive disorder NOS
- 29/41 cases of cognitive disorder were seen in low titer patients
- Seizures were seen in 10/236 (4%), 7 in low titer patients
- Cerebellar ataxia in 9/236 (3.8%), 5 in low titer patients
- Headache disorders in 9/236 (3.8%)
- 30% of patients had co-existing neural autoantibodies
- CSF abnormal in 81%; elevated protein in 55 (23%), leukocytosis in 13 (5%)
- Authors recommend that seropositive results be interpreted in the clinical context of the individual patient

Zalewski et al. 2016



## VGCC ab Encephalitis

- Auto-antibodies against P/Q- and N-type voltage dependent calcium channels mimicking frontotemporal dementia
- 54 year old man presented with frontal disinhibition and cognitive decline over days to weeks; had a history of rheumatoid arthritis and thyroid eye disease
- MRI showed atrophy greater than would be expected for age; FDG-PET consistent with FTD
- Symptoms initially treated with medication but symptoms worsened and paraneoplastic panel obtained due to patient's history of autoimmune conditions
- Elevated antibodies to VGCC abs +0.24 (seropositive medium)
- Patient underwent 5 therapeutic plasma exchange sessions; notable improvement starting with third session

Younes et al. 2018



## Encephalopathies & Dementias

- Autoimmune dementia and encephalopathies are complex disorders that can cause immune-mediated cognitive deficits and have confusing nomenclature
- Presentation varies from acute limbic encephalitis to subacute or chronic disorders of cognition mimicking neurodegenerative dementia
- May occur as a paraneoplastic phenomenon or an idiopathic autoimmune phenomenon.
- Important to evaluate for cancer
- The diagnosis of an autoimmune dementia requires the detection of objective improvements in cognitive decline after a course of immunotherapy

McKeon et al. 2020; McKeon 2016; Flanagan et al. 2026



## Patient Background Information

- 54 yr old right-handed SM (Service Member) born and raised in Nigeria
- Grew up speaking both English and his native language
- Completed school through university in Nigeria, Master's Degree in Leadership Management through Liberty University while on active duty
- Achieved the rank of MSG (Master Sergeant, E-8), Military Occupational Specialty of 36B, Financial Management Technician
- 19 years time in service
- Married for 17 years, 5 children
- One deployment to Iraq, November 2005-November 2006; history of fall from helicopter 2005, blast exposure 2006 with no noted residual sequelae.
- Wife noted symptoms of PTSD upon SM's return from deployment



## History of the Presenting Problem

- SM and wife abducted and held for 5 days while on a trip to visit family in Nigeria in December 2015
- SM reportedly beaten to include several blows to the head
- Diagnosed with Major Depressive Disorder in February 2016
- Very prominent psychomotor slowing, flat affect, looking at the floor and having difficulty making eye contact
- TBI Team noted mild confusion, poor attention/concentration, word-finding problems, difficulty getting to work on time, frequent HA's, occasionally hearing voices
- Commander concerned about "very slow to respond"
- Brief NP testing indicated general cognitive performance in the extremely low range of functioning
- Examiner concluded results were not indicative of SM's true neuropsychological status, multiple failures on PVT's



## History of the Presenting Problem

- Neurology exam in May 2016 found incongruous R superior quadrantanopia
- MRI June 2016 scattered focal areas of R greater than L bilateral volume loss in the cerebellar hemispheres
- Also very minimal smooth diffuse pachymeningeal enhancement, more prominent in the anterior temporal lobes and overlying parietal convexities, could be wnl but radiologist suggested correlation with signs and symptoms of infectious etiology
- SM sent for inpatient treatment for PTSD and MDD in San Antonio
- Referred for NP testing by psychiatrist, degree of cognitive impairment seemed out of proportion for even severe PTSD and MDD



## Past Medical & Symptom History

- AHLTA Electronic Medical Record (EMR)
- Collateral Interview with wife
- Wife noted problems with memory beginning in 2010, forgot children's birthdates for paperwork, called home to ask
- History of HA dating to 2010, diagnosed as migraine HA in March 2012 after a HA that lasted 2-3 days
- MRI in 2010 a small nonenhancing area of abnormal signal in the posterior medial right cerebellar hemisphere thought to suggest infarction at that time
- Additional medical problems in 2010 included chronic leukopenia, hx of pulmonary embolism, hx deep vein thrombosis x2, erectile dysfunction (autonomic? High percent in LEMS) and chronic back pain



## Past Medical & Symptom History

- 2012 Memory deficits brought to the SM's attention by co-workers
- **November 2012 Neurology exam: deficits in recall, multi-step commands, and visuospatial testing; twice incorrectly gave the year as 2011**
- NP evaluation ordered but was not completed at that time
- MRI 2012 reported mixed porencephaly and encephalomalacia involving the right cerebellar hemisphere
- Noted in 2012 several month history of anxiety, depression, insomnia
- SM diagnosed with PTSD in 2014
- Per wife's report, SM had experienced auditory hallucinations for several years, prior to abduction, could not be attributed to severe PTSD



## Presentation & NP Testing

- Seen for NP testing August 2016
- Upon presentation to this provider the SM demonstrated very prominent psychomotor slowing, flat affect with little eye contact, speech was very sparse, mild occasional stuttering noted, SM was able to communicate that he was having difficulty with word-finding
- Neurocognitive functioning severely and almost globally impaired
- FSIQ=50
- NP attempted Category Test, 14 minutes to complete first two categories, D/C before Category III
- NP administered WCST, 55 minutes to administer, strongly perseverated to Form for about half of the test, then more disorganized, might get 3-5 correct but 4 FMS
- Trails B allowed to finish, 8+ minutes



## Presentation & NP Testing


- CVLT-2 Moderate to severe impairment but able to give main points of the 2 LM passages and at delayed recall as well
- Visual Reproduction recalled a few bits of information and retained about half of those at 30 minutes
- Surprisingly copy of ROCF was accurate and well-organized, lost organization at immediate and delayed recall
- Performance in severely impaired range bilaterally on Finger Tapping, consistent with very slowed presentation
- Failed all PVT's
- SM appeared able to read and respond to BDI and BAI, reported clinically significant levels of depression and anxiety but no SI
- Wife reported declining levels of cognitive function since 2010, at time of evaluation unable to wash dishes correctly; ADL's intact but needed reminding; IADL's very impaired; got lost if not accompanied






## Recommendations & Followup

- NP diagnosis of Major Neurocognitive Disorder (dementia)
- Recommended further evaluation for etiology of dementia, to include any possible reversible causes, specifically requested Neurology rule out limbic encephalitis
- Referred to Neuro-ophthalmology due to previous finding of visual field defects
- EEG September 2017 Clinical Impression: voltage of the electrocerebral activity very low, including the posterior dominant rhythm, but essentially normal
- Neuro-ophthalmology found bilateral superior visual field defects, but again repeat MRI did not find correlated lesions for visual field defects
- MRI findings January 2017 read as stable R posterior inferior cerebellar encephalomalacia compared to June 2016 MRI



## Coronal T1w/contrast




## Neurology

- Neurologist thought findings suggestive of autoimmune encephalitis: past finding of protein in CSF 50% higher than normal; diffuse pachymeningeal enhancement; chronic HA's 3-4 times per week
- FDG PET found no evidence for FTD
- Genetic testing for genetic variants associated with FTD and ALS negative
- Paraneoplastic panel positive for N-type calcium channel antibodies
- Low clinical range 0.04-0.09 nmol/L, pt's results 0.07
- Neurologist felt nonparaneoplastic more likely due to long time course of decline
- Thus far continued evaluation for cancer to include full body PET negative, but some suspicious prostate findings
- Inpatient treatment with IVIG for 4 days, followed by treatment with monthly IVIG has produced functional improvement, thus far no repeat NP testing



## Summary

- This case study illustrates the course of a progressive dementia over about a 5 year period of time, seropositive for VGCC N-type antibody
- Some recovery with treatment; potential for recovery unclear
- Consideration of an autoimmune encephalitis as one of the possible reversible causes for progressive dementia may result in earlier diagnosis, earlier treatment, better recovery







## References

- Annunziata, G., Lobo, P., & Carbuccia, C. (2017) A rare case of cerebellar ataxia due to voltage-gated calcium channel and glutamic acid decarboxylase autoantibodies. *Am J. Case Rep* 2017 Nov 27, 18: 1251-1255.
- Bekircan-Kurt, C., Ciftci, E., Kurme, A., & Anlar, B. (2015) Voltage gated calcium channel anti-body-related neurological diseases. *World Journal of Clinical Cases*, 3(3), 293-300.
- Dalmau, J., DeAngelis, L.M., & Eichler, A.F. (July 2018). Paraneoplastic and autoimmune encephalitis. UpToDate. [www.uptodate.com](http://www.uptodate.com)
- Finke, C., Kopp, U., Pruss, H., Dalmau, J., Wandinger, K-P & Pioner, C. (2012) Cognitive deficits following anti-NMDA receptor encephalitis. *J Neurol Neurosurg Psychiatry*, 83 (2): 195-198.
- Finkel, L. & Kon, S. (2013). N-type calcium channel antibody-mediated autoimmune encephalitis: An unlikely cause of a common presentation. *Epilepsy Behav Case Rep*, 1: 92-96. (case of a 14-year old boy)
- Flanagan, E., Drubach, D., & Boeve, B. (2016). *Handb Clin Neurol* 133: 247-67.
- Lancaster, E. (2016) The diagnosis and treatment of autoimmune encephalitis. *J Clin Neurol* 12(1) 1-13.



## References

- Loughan, A., Allen, A., Perna, R., & Malkin, M. (2016). Anti-N-Methyl-D-Aspartate Receptor Encephalitis: A review and neuropsychological case study. *The Clinical Neuropsychologist*, 30 (1), 150-163.
- McKeon, A., Lennon, V.A., & Pittock, (2016) Autoimmune Encephalopathies and Dementias. *Continuum (Minneapolis, Minn)* Apr;24(2 Dementia) 538-58.
- McKeon, A. (2013) Paraneoplastic and other autoimmune disorders of the central nervous system. *Neurohospitalist*, 3(2) 53-64.
- McKeon, A., Lennon, V.A., & Pittock, (2010) Immunotherapy-responsive dementias and encephalopathies. *Continuum (Minneapolis, Minn)* Apr;16(2 Dementia) 80-101.
- McKeon, A., Marnane, M., O'Connell, M., Stack, J., Kelly, P. & Lynch, T. (2007) Potassium channel antibody associated encephalopathy presenting with a frontotemporal dementia like syndrome. *Archives of Neurology* 64 (10), 1528-1530.
- Tuzun, E. & Dalmau, J. (2007) Limbic encephalitis and variants: Classification, diagnosis and treatment. *The Neurologist*, 13: 365-374.
- Younes, K., Lepow, L, Estrada, C., & Schultz, P. (2018). Auto-antibodies against P/Q and N-type voltage-dependent calcium channels mimicking frontotemporal dementia. *SAGE Open Med Case Report*. Published online April 10, 2018.
- Zaleski, N., Lennon, V., Lachance, D., Klein, C., Pittock, S., & McKeon, A. (2016) P/Q- & N-type calcium channel antibodies: Oncological, neurological & serological accompaniments. *Muscle Nerve* 54 (2): 220-227.